

# Multidimensional analogues of Bohr's theorem on power series\*

Lev Aizenberg

March 30, 1998

## Abstract

Generalizing the classical result of Bohr, we show that if an  $n$ -variable power series converges in an  $n$ -circular bounded complete domain  $D$  and its sum has modulus less than 1, then the sum of the maximum of the moduli of the terms is less than 1 in the homothetic domain  $r \cdot D$ , where  $r = 1 - \sqrt[n]{2/3}$ . This constant is near to the best one for the domain  $D = \{z : |z_1| + \dots + |z_n| < 1\}$ .

## 1 Preliminaries

It is known the following classical Bohr's result [5], which final form was contributed by M. Riesz, I. Schur and N. Wiener.

**Theorem 1** *If a power series*

$$\sum_{k=0}^{\infty} c_k z^k \tag{1}$$

*converges in the unit disk and its sum has modulus less than 1 then*

$$\sum_{k=0}^{\infty} |c_k z^k| < 1 \tag{2}$$

*in the disk  $\{z : |z| < 1/3\}$  and the constant  $1/3$  cannot be improved.*

---

\*Middle East Technical University Preprint Mathematics 164/98. This work was supported by the BSF grant No 94-00113.

Recently H. P. Boas and D. Khavinson obtained some multidimensional generalization of this result ([4], see also for more references there). Denote by  $K_n$  the largest number such that if the series

$$\sum_{\alpha} c_{\alpha} z^{\alpha} \quad (3)$$

converges in the unit polydisk  $U_1 = \{z : |z_j| < 1, j = 1, \dots, n\}$  and the estimate

$$\left| \sum_{\alpha} c_{\alpha} z^{\alpha} \right| < 1, \quad (4)$$

is valid there, then

$$\sum_{\alpha} |c_{\alpha} z^{\alpha}| < 1, \quad (5)$$

holds in  $K_n \cdot U_1$ ; here  $\alpha = (\alpha_1, \dots, \alpha_n)$ , all  $\alpha_j$  are non-negative integers,  $z = (z_1, \dots, z_n)$ ,  $z^{\alpha} = z_1^{\alpha_1} \dots z_n^{\alpha_n}$ .

**Theorem 2 (Boas, Khavinson)** *It is true for  $n > 1$  that*

$$\frac{1}{3\sqrt[n]{n}} < K_n < \frac{2\sqrt{\log n}}{\sqrt[n]{n}} \quad (6)$$

**Theorem 3 (Boas, Khavinson)** *Let the series (3) converge in a complete  $n$ -circular domain (Reinhardt's domain)  $D$  and (4) holds in  $D$ . Then (5) is true in the homothetic domain  $K_n \cdot D$ .*

Notice that Remark 1 from [4], in fact, contains a result stronger than the left part of the inequality (6), namely

$$K_n > \begin{cases} \frac{2}{5\sqrt[n]{n}} & \text{for } n > 1, \\ \frac{1}{2\sqrt[n]{n}} & \text{for large enough } n. \end{cases}$$

## 2 Main Results

We consider some other multidimensional variants of Bohr's problem. Denote by  $B_n(D)$  the biggest number  $r$  such that if the series (3) converges in a complete  $n$ -circular domain  $D$  and (4) holds in it then

$$\sum_{\alpha} \sup_{D_r} |c_{\alpha} z^{\alpha}| < 1, \quad (7)$$

where  $D_r = r \cdot D$  is the homothetic transformation of  $D$ . If  $D = U_1$  then  $B_n(D) = K_n$ . In particular, our consideration is also a natural generalization of Bohr's theorem (Theorem 1). Notice that it is shown in [2] that any power series (3), converging in  $D$ , converges in the sense of the left part of (7) for all  $r$ ,  $0 < r < 1$ .

**Theorem 4** *The inequality*

$$1 - \sqrt[n]{\frac{2}{3}} < B_n(D). \quad (8)$$

*is true for any complete bounded  $n$ -circular domain  $D$ .*

This estimate can be improved for concrete domains.

**Theorem 5** *For the unit ball  $D^1 = \{z : |z_1|^2 + \dots + |z_n|^2 < 1\}$  the following estimate is true:*

$$B_n(D^1) > \begin{cases} \frac{2}{5n} & \text{for } n > 1, \\ \frac{1}{2n} & \text{for large enough } n. \end{cases}$$

**Theorem 6** *For the unit hypercone  $D^0 = \{z : |z_1| + \dots + |z_n| < 1\}$  the following inequality holds:*

$$B_n(D^0) < \frac{0,446663}{n}. \quad (9)$$

**Corollary.**

$$1 - \sqrt[n]{\frac{2}{3}} < B_n(D^0) < \frac{0.446663}{n}. \quad (10)$$

**Remark 1.** The following asymptotic equality

$$1 - \sqrt[n]{\frac{2}{3}} = \frac{\log 3/2}{n} + O\left(\frac{1}{n^2}\right)$$

is true, where  $\log 3/2 \approx 0.405465$ . Denoting by  $B_-$  the left part of (10) and by  $B_+$  the right one, we get

$$1 < \lim_{n \rightarrow \infty} \sup \frac{B_+}{B_-} < 1,1016.$$

Consider now another problem. Denote by  $L_n(D^\circ)$  the biggest number  $r$  such that if the series (3) converges in the hypercone  $D^\circ$  and (4) holds in it, then

$$\sum_{\alpha} \|c_{\alpha} z^{\alpha}\|_{L_1(\partial D_r^\circ)} < 1, \quad (11)$$

where the  $L^1$ -norm is considered with respect to the measure  $\mu_r$ , which is the image of the measure

$$d\mu = \frac{(n-1)!}{(2\pi i)^n} d|z_1| \wedge \cdots \wedge d|z_{n-1}| \wedge \frac{dz_1}{z_1} \wedge \cdots \wedge \frac{dz_n}{z_n}$$

by the homothetic transformation  $z \rightarrow rz$ . Usually the measure  $d\mu$  is used for calculating of Szego kernel for  $D^0$  (see [3]); notice that  $\mu(\partial D^0) = 1$ . For  $n = 1$  this problem coincides with Bohr's problem. The analogous problem for the polydisk  $U_1$  (with  $L^1$  on its Silov boundary with respect to usual Lebesgue measure on it) is equivalent to the problem, considered in Theorem 2.

**Theorem 7** *For the hypercone  $D^0$  the following estimates are true:*

$$0,238843 < L_n(D^\circ) \leq 1/3. \quad (12)$$

It is surprising here that the estimates in (12) does not depend on  $n$ , which is different from the results of theorems 2—6.

Now consider a variant of multidimensional Bohr's problem, dealing with expansions in series of homogeneous polynomials, which is also a natural generalization power series expansion. Let  $Q$  be a complete circular domain (Cartan's domain) centered at  $0 \in Q$ ; then any function  $f(z)$ , holomorphic in  $Q$  can be expanded into the series

$$f(z) = \sum_{k=0}^{\infty} P_k(z), \quad (13)$$

where  $P_k(z)$  is a homogeneous polynomial of degree  $k$ .

**Theorem 8** *If the series (13) converges in the domain  $Q$  and the estimate  $|f(z)| < 1$  holds in it, then*

$$\sum_{k=0}^{\infty} |P_k(z)| < 1 \quad (14)$$

*in the homothetic domain  $(1/3)Q$ . Moreover, if  $Q$  is convex, then  $1/3$  is the best possible constant.*

### 3 Proofs

**Proof of Theorem 4.** The following generalization of Cauchy inequalities was considered in [2]: if (4) holds in  $D$  then

$$|c_\alpha| < \frac{1}{d_\alpha(D)}, \quad (15)$$

where  $d_\alpha(D) = \max_{\bar{D}} |z^\alpha|$ . Using Wiener's method it is easy to strengthen the estimates (15):

$$|c_\alpha| < (1 - |c_0|^2) \frac{1}{d_\alpha(D)} \quad (16)$$

for  $|\alpha| = \alpha_1 + \dots + \alpha_n > 1$ . We do not present the proof of (16), because it repeats the proof for polydisk  $U_1$  (see [4]) but deals with inequalities (15) instead of Cauchy inequalities. If (4) holds in  $D$  then, applying (16), we get

$$\begin{aligned} \sum_{\alpha} \sup_{D_r} |c_\alpha z^\alpha| &= \sum_{\alpha} |c_\alpha| d_\alpha(D_r) < |c_0| + \\ &+ (1 - |c_0|^2) \sum_{|\alpha|=1}^{\infty} \frac{d_\alpha(D_r)}{d_\alpha(D)} = |c_0| + (1 - |c_0|^2) \sum_{k=1}^{\infty} \binom{n+k-1}{k} r^k = \\ &= |c_0| + (1 - |c_0|^2) \left[ \frac{1}{(1-r)^n} - 1 \right]. \end{aligned}$$

If now

$$\frac{1}{(1-r)^n} - 1 \leq \frac{1}{2}, \quad (17)$$

then

$$\sum_{\alpha} \sup_{D_r} |c_\alpha z^\alpha| < |c_0| + (1 - |c_0|^2) \frac{1}{2} = 1 - \frac{1}{2}(1 - |c_0|)^2 < 1.$$

The condition (17) means that (7) is true if  $r \leq 1 - \sqrt[n]{\frac{2}{3}}$ .

**Proof of Theorem 5.** Consider Borel probability measure on  $\partial D^1$ , which is invariant under all unitary transformations of  $\mathbf{C}^n$ :

$$d\mu = \frac{(n-1)!}{(2\pi i)^n} d|z_1|^2 \wedge \dots \wedge d|z_{n-1}|^2 \wedge \frac{dz_1}{z_1} \wedge \dots \wedge \frac{dz_n}{z_n}.$$

The monomials  $z^\alpha$  are orthogonal with respect to integration by  $\mu$  and

$$\int_{\partial D^1} |z^{2\alpha}| d\mu = \frac{\alpha_1! \dots \alpha_n! (n-1)!}{(|\alpha| + n - 1)!}. \quad (18)$$

Further we repeat the proof of Theorem 2 from [4] by Wiener method, but integrating on the sphere  $\partial D^1$  by the measure  $\mu$  instead of integrating on the unit torus as in [4]. We obtain

$$\sum_{|\alpha|=k} |c_\alpha|^2 \frac{\alpha_n! \cdots \alpha_n! (n-1)!}{(|\alpha| + n - 1)!} < (1 - |c_0|^2)^2.$$

Furthermore, notice that from Schwarz Lemma for the ball  $D^1$  it follows (again by using Wiener's method from [4])

$$\sum_{|\alpha|=1} |c_\alpha|^2 < (1 - |c_0|^2)^2.$$

Then, remembering that

$$d_\alpha(D^1) = \sqrt{\frac{\alpha_1^{\alpha_1} \cdots \alpha_n^{\alpha_n}}{|\alpha|^{|\alpha|}}},$$

where  $0^0 = 1$ , we get

$$\begin{aligned} & \sum_{\alpha} \sup_{D_r} |c_\alpha z^\alpha| = |c_0| + \left( \sum_{|\alpha|=1} |c_\alpha| \right) r + \\ & + \sum_{k=2}^{\infty} \sum_{|\alpha|=k} |c_\alpha| \sqrt{\frac{\alpha_1^{\alpha_1} \cdots \alpha_n^{\alpha_n}}{|\alpha|^{|\alpha|}}} r^k < \\ & < |c_0| + (1 - |c_0|^2) r \sqrt{n} + (1 - |c_0|^2) \sum_{k=2}^{\infty} \left( \sum_{|\alpha|=k} \frac{(k+n-1)! \alpha_1^{\alpha_1} \cdots \alpha_n^{\alpha_n}}{k^k \alpha_1! \cdots \alpha_n! (n-1)!} \right)^{1/2} r^k < \\ & < |c_0| + (1 - |c_0|^2) r \sqrt{n} + (1 - |c_0|^2) \sum_{k=2}^{\infty} \left( \frac{(k+n-1)!}{k! (n-1)!} \sum_{|\alpha|=k} \frac{k!}{\alpha_1! \cdots \alpha_n!} \right)^{1/2} r^k = \\ & = |c_0| + (1 - |c_0|^2) r \sqrt{n} + (1 - |c_0|^2) \sum_{k=2}^{\infty} \sqrt{\binom{k+n-1}{k}} (r \sqrt{n})^k \end{aligned}$$

It remains now to apply the estimate, obtained in Remark 1 from [4].

**Proof of Theorem 6.** Consider the function

$$f_a(z) = \frac{1+a}{2} \frac{1 - (z_1 + \cdots + z_n)}{1 - a(z_1 + \cdots + z_n)} = \sum_{\alpha} c_\alpha z^\alpha,$$

where  $0 < a < 1$ ; then  $|f_a(z)| < 1$  in hypercone  $D^\circ$  and

$$\sum_{\alpha} |c_{\alpha} z^{\alpha}| = \frac{1+a}{2} + \frac{1-a^2}{2} \sum_{k=1}^{\infty} a^{k-1} \sum_{|\alpha|=k} \frac{k!}{\alpha_1! \cdots \alpha_n!} |z^{\alpha}|.$$

Since

$$d_{\alpha}(D^\circ) = \frac{\alpha_1^{\alpha_1} \cdots \alpha_n^{\alpha_n}}{|\alpha|^{|\alpha|}},$$

it follows

$$\begin{aligned} \sum_{\alpha} |c_{\alpha}| d_{\alpha}(D_r^\circ) &= \frac{1+a}{2} + \frac{1-a^2}{2} \sum_{k=1}^{\infty} \sum_{|\alpha|=k} a^{k-1} \frac{k! \alpha_1^{\alpha_1} \cdots \alpha_n^{\alpha_n}}{\alpha_1! \cdots \alpha_n! k^k} r^k > \\ &\geq \frac{1+a}{2} + \frac{1-a^2}{2} \sum_{k=1}^{\infty} a^{k-1} \frac{k!}{k^k} \left( \sum_{|\alpha|=k} \frac{1}{\alpha_1! \cdots \alpha_n!} \right) r^k = \\ &= \frac{1+a}{2} + \frac{1-a^2}{2a} \sum_{k=1}^{\infty} \frac{(anr)^k}{k^k} \geq 1, \end{aligned}$$

if

$$\sum_{k=1}^{\infty} \frac{(anr)^k}{k^k} \geq \frac{a}{1+a}.$$

Let  $x_0(a)$  be the root of the equation

$$\sum_{k=1}^{\infty} \frac{x^k}{k^k} = \frac{a}{1+a};$$

then, if  $anr \geq x_0(a)$ , (7) fails for that  $r$  and  $D = D^0$ . Now considering  $a \rightarrow 1$ , we obtain that (7) is not true for  $D = D_0$  if  $r \geq \frac{x_0}{n}$ , where  $x_0 = x_0(1)$ . Take notice that  $x_0$  is a root of the equation

$$\sum_{k=1}^{\infty} \frac{x^k}{k^k} = 1/2, \tag{19}$$

hence  $B_n(D^0) \leq x_0/n$ . By using the program “Mathematica 3.0” [7], we estimated from above  $x_0$ . We obtained that the equation

$$\sum_{k=1}^p \frac{x^k}{k^k} = 1/2 \tag{20}$$

has a root 0,446662 (where the last decimal digit is precise) if  $p$  runs from 5 till 25. So the equation (20) has a root less than 0,446663 if  $5 \leq p \leq 25$ , hence this estimate is true for  $x_0$ .

**Proof of Theorem 7.** Take notice that by analogy with (18) it is true that

$$\int_{\partial D_r^\circ} |z^\alpha| d\mu_r = \frac{\alpha_1! \cdots \alpha_n! (n-1)!}{(|\alpha| + n - 1)!} r^{|\alpha|}.$$

Using (16), we get

$$\begin{aligned} \sum_{\alpha} \|c_{\alpha} z^{\alpha}\|_{L^1(\partial D_r^\circ)} &< |c_0| + (1 - |c_0|^2) \sum_{k=1}^{\infty} \sum_{|\alpha|=k} \\ &\frac{k^k (n-1)!}{(n+k-1)!} \frac{\alpha_1! \cdots \alpha_n!}{\alpha_1^{\alpha_1} \cdots \alpha_n^{\alpha_n}} r^{|\alpha|} < |c_0| + (1 - |c_0|^2) \times \\ &\times \sum_{k=1}^{\infty} \frac{k^k (n-1)!}{(n+k-1)!} \sum_{|\alpha|=k} r^{|\alpha|} = |c_0| + (1 - |c_0|^2) \sum_{k=1}^{\infty} \frac{k^k}{k!} r^k. \end{aligned}$$

Denote by  $r_0$  the root of the equation

$$\sum_{k=1}^{\infty} \frac{k^k}{k!} r^k = \frac{1}{2}, \quad (21)$$

then (11) holds for  $r = r_0$ , therefore  $L_n(D^0) \geq r_0$ . The left side of (21) is less than the left side of the equation

$$\sum_{k=1}^{25} \frac{k^k}{k!} r^k + \frac{1}{\sqrt{52\pi}} \frac{(ex)^{26}}{1 - ex} = 1/2, \quad (22)$$

what follows from an approximation of Gamma-function. Using again “Mathematica 3.0”, we show that the root of the equation (22) is bigger than 0,238843, therefore the left part of the inequality (12) is true.

Further, consider, as in proof of Theorem 6, a function  $f_a(z)$ , for which

$$\begin{aligned} \sum_{\alpha} \|c_{\alpha} z^{\alpha}\|_{L^1(\partial D_r^\circ)} &= \frac{1+a}{2} + \frac{1-a^2}{2} \sum_{k=1}^{\infty} \frac{k!(n-1)!a^{k-1}}{(n+k-1)!} \times \\ &\times \sum_{|\alpha|=k} r^{|\alpha|} = \frac{1+a}{2} + \frac{1-a^2}{2a} \sum_{k=1}^{\infty} (ar)^k \geq 1, \end{aligned}$$

if

$$\sum_{k=1}^{\infty} (ar)^k \geq \frac{a}{1+a}.$$

When  $a \rightarrow 1$ , we get that the inequality (11) fails if  $r > 1/3$ .

**Proof of Theorem 8.** In each section of the domain  $Q$  by a complex line

$$\alpha = \{z : z_j = a_j t, \quad j = 1, \dots, n; \quad t \in \mathbf{C}\}$$

the series turns into the power series by  $t$

$$f(at) = \sum_{k=0}^{\infty} P_k(a) t^k$$

and, in addition,  $|f(at)| < 1$ . By Theorem 1

$$\sum_{k=0}^{\infty} |P_k(a) t^k| < 1$$

in the section  $\alpha \cap (\frac{1}{3} \cdot Q)$ . But it is just (14), since  $\alpha$  is an arbitrary complex line passing through the origin. Inversely, let the domain  $Q$  be convex, then  $Q$  is an intersection of half-spaces

$$Q = \bigcap_{a \in J} \{z : \operatorname{Re}(a_1 z_1 + \dots + a_n z_n) < 1\}$$

with some  $J$ . Since  $Q$  is circular, we obtain

$$Q = \bigcap_{a \in J} \{z : |a_1 z_1 + \dots + a_n z_n| < 1\}.$$

It is sufficient now to show that the constant  $1/3$  cannot be improved for each domain  $P_a = \{z : |a_1 z_1 + \dots + a_n z_n| < 1\}$ . From Theorem 1 it follows that for any  $r > 1/3$  there exists a function  $f(z)$ , represented by (1) and such that  $|f(z)| < 1$  in the unit disk, but (2) fails in the disk  $\{z : |z| < r\}$ . To finish the proof we use the functions  $f(a_1 z_1 + \dots + a_n z_n)$ .

## 4 Final Remarks

**Remark 2.** In the proof of Theorem 5 the fact, that the domain in consideration is the ball  $D^1$  is used twice as follows: in order to prove that the monomials  $z^\alpha$  are orthogonal on the sphere  $\partial D^1$  with respect to the measure  $d\mu$  and to get the estimate  $d_\alpha(D^1) \leq 1$ . Therefore an analogous theorem holds for any complete  $n$ -circular domain  $D$  if  $D^1 \subset D \subset U_1$ . But, for the hypercone  $D^\circ$  this is not true (see Theorem 6).

**Remark 3.** In the proof of Theorem 6 was used the following quite rough inequality:

$$\sum_{|\alpha|=k} \frac{\alpha_1^{\alpha_1} \cdots \alpha_n^{\alpha_n}}{\alpha_1! \cdots \alpha_n!} \geq \sum_{|\alpha|=k} \frac{1}{\alpha_1! \cdots \alpha_n!}.$$

Therefore, in fact, the estimate from above in the theorem might be decreased. For example, if  $n = 2$  (10) implies  $B_2(D^0) < 0.223332$ , but using PC one can show that  $B_2(D^0) < 0.191373$ . Hence,  $0.183502 < B_2(D^\circ) < 0.191373$ .

**Remark 4.** Comparing Theorem 2 and Corollary we can see that  $B_n(D)$  depends essentially on the domain  $D$  since  $B_n(U_1) = K_n$ . It seems that it is more natural to consider not a single number in the problem, considered in Theorem 2 and Theorem 3, but the largest subdomain  $D_B$  of  $D$  such that (5) holds. From [4] it follows, for example, that  $U_{1B}$  contains the ball  $\frac{1}{3} \cdot D^1$ .

**Remark 5.** Notice that there exist such unbounded  $n$ -circular domains  $D$  for which the problems defined by the conditions (5) and (7), are equivalent. Consider, for example, the domains  $D = \{z : |z|^\beta < c\}$ , where  $\beta$  is a multi-index with coprime components. Bounded holomorphic functions in such domains depend only on one variable  $z^\beta$  and the exact value of Bohr's radius equals to  $(\frac{1}{3})^{1/|\beta|}$ . Thus there exists  $n$ -circular domains with Bohr's radius arbitrarily close to 1. Therefore it is impossible to remove the assumption about convexity in Theorem 8.

**Remark 6.** Unlike Theorems 1—7, in Theorem 8 the series (13) is not a basis expansion. In [1] it was shown that there exists basis in the space of all holomorphic functions in  $Q$ , consisting of homogeneous polynomials  $P_{k,m}(z)$ , where  $k$  is the degree of the polynomial and  $m = 1, \dots, \binom{k+n-1}{k}$ . It is reasonable to consider Bohr's problem for such basis expansions, but there is no results yet. This question is a particular case of a more general problem in Bohr spirit for expansions by an arbitrary basis in a domain  $D \subset \mathbf{C}^n$  (under some restrictions on the basis, because, for example, the basis  $\{1, (z-1)/2, z^2, z^3, \dots\}$  in the unit disk has no Bohr's constant).

## Acknowledgment

I am sincerely thankful to D. Khavinson, who turned my attention to Bohr's result (Theorem 1) and its first multidimensional analogues (Theorems 2 and 3), and to V. Zahariuta and P. Djakov for their help in preparing the English version of the paper.

The results of this paper were obtained during the author's visit in the Middle East Technical University. The author would like to thank A. Aytuna and S. Alpay for the invitation, and as well, A. Aytuna and B. Karasözen for their support.

## References

- [1] L.A. Aizenberg, The spaces of functions analytic in  $(p, q)$ -circular regions, *Sov. Math. Dokl.* **2** (1960), 79-82.
- [2] L.A. Aizenberg, B.S. Mityagin, The spaces of functions analytic in multicircular domains, *Sibirs. Math. J.* **1** (1960), 153-170 (Russian).
- [3] L.A. Aizenberg, A.P. Yuzhakov, *Integral Representations and Residues in Multidimensional Complex Analysis*, AMS, 1983, 283 pp.
- [4] H.P. Boas, D. Khavinson, Bohr's power series theorem in several variables, *Proc. Amer. Math. Soc.* **125** (1997), 2975-2979.
- [5] H. Bohr, A theorem converging power series, *Proc. London Math. Soc.* (2) **13** (1914) 1-5.
- [6] W. Rudin, *Function theory in the unit ball of  $\mathbf{C}^n$* , Springer-Verlag, 1980, 436 pp.
- [7] Wolfram Research, *Mathematica 3.0*, 1996.

Address:

Department of Mathematics  
and Computer Science,  
Bar-Ilan University  
52900 Ramat-Gan, ISRAEL  
e-mail: aizenbrg@macs.biu.ac.il

Temporary address:

Department of Mathematics  
Middle East Technical University  
06531 Ankara, TURKEY  
e-mail: aizenbrg@arf.math.metu.edu.tr